

Therapeutic approaches in arterial thrombosis

D. R. PHILLIPS, P. B. CONLEY, U. SINHA and P. ANDRE

Portola Pharmaceuticals, Inc., San Francisco, CA, USA

To cite this article: Phillips DR, Conley PB, Sinha U, Andre P. Therapeutic approaches in arterial thrombosis. *J Thromb Haemost* 2005; 3: 1577–89.

Summary. The current standard of care for the treatment of arterial thrombosis includes anticoagulants and three classes of antiplatelet agents – aspirin, thienopyridines and glycoprotein IIb-IIIa antagonists. Although these drugs have had a significant impact on morbidity and mortality in several patient populations, up to 15% of the high risk patients with acute coronary syndrome continue to suffer from ischemic events. This problem may occur, in part, because the platelets in many patients are non-responsive to aspirin and clopidogrel. Murine models now indicate that platelets are not only responsible for arterial occlusion, they are also involved in the progression of atherosclerotic disease. New opportunities have emerged identifying potential targets and strategies for drug discovery suited to address these deficiencies by more effectively modulating platelet adhesion, thrombus growth, thrombus stability and the pro-inflammatory activity of platelets. In addition, a growing need has emerged for the development of bedside devices capable of bringing personalized medicine to patients being treated with antithrombotic drugs in order to measure the pharmacodynamic activities of new therapies, to assess the activities achieved by combined antithrombotic therapy, and to identify patients that fail to respond.

Keywords: anticoagulant, antiplatelet, atherosclerosis, platelet, platelet monitoring, thrombosis.

Introduction

Arterial thrombosis is the result of sequential events involving platelet adhesion, activation and subsequent aggregation that can lead to vascular occlusion, perhaps the primary pathological complication of advanced atherosclerotic lesions. Recent advances in the field of thrombosis suggest that the second pathological consequence of platelet adhesion and activation may be as consequential as the immediate ischemia induced by arterial thrombosis as platelets are a primary source of several inflammatory proteins known to be involved in the progression of atherosclerotic disease including RANTES, sCD40L, PDGF and transforming growth factor- β (TGF- β). These considerations suggest that therapeutic targeting of platelets

has two objectives: first, prevention of vessel occlusion; second, inhibition of the platelet contribution to lesion progression.

The pharmaceutical industry has made important inroads into the development of drugs for the treatment of the thrombotic complications of atherosclerosis. In one example, the occlusive, ischemic consequences of acute myocardial infarction (MI) have been addressed, by thrombolytics to lyse the thrombus, and more recently by interventional strategies to mechanically remove or dislodge the thrombus and to maintain artery patency with stents coated with agents such as rapamycin or paclitaxel to reduce the incidence of restenosis. In yet another example, antagonists against platelet receptors such as glycoprotein (GP) IIb-IIIa and P2Y₁₂ have been developed, joining aspirin, a Cox-1 inhibitor as the primary antithrombotic drugs. However, despite these advances in antithrombotic therapies and the widespread use of statins to reduce cholesterol and CRP levels, the incidence of atherosclerosis continues to rise, as do the ischemic consequences of atherosclerosis including MI and stroke. An added complication is type 2 diabetes which is an independent risk factor for cardiovascular disease (CVD). A recent analysis of the Framingham Heart Study showed that even though management of risk factors such as blood pressure and cholesterol has improved significantly in the total patient population, the presence of diabetes significantly reduced the overall benefits [1]. While it may be possible to address these pathologies by more aggressive health management, and by more optimal application of existing therapies, clearly, truly effective treatment of the thrombotic consequences of atherosclerosis requires not only the development of drugs to be used as primary care on a chronic basis to prevent thrombosis and its ischemic complications but also to block the contribution of platelet-induced inflammation in the progression of atherosclerotic disease. Reviewed below is the mechanism of action, clinical successes and limitations of the four drug classes currently used to prevent arterial thrombosis; aspirin, P2Y₁₂ inhibitors, GP IIb-IIIa antagonists and anticoagulants. Therapeutic opportunities afforded by our current understanding of the mechanisms of arterial thrombosis and the inflammatory activity of platelets are discussed. Finally, recognizing that individuals vary in response to various drugs and that combination antithrombotic therapies has become commonplace, we will highlight the need for improved pharmacodynamic assessment of platelet function.

Correspondence: David R. Phillips, Portola Pharmaceuticals, Inc., 270 E. Grand Ave., South San Francisco, CA 94402, USA.
Tel.: +1 650 246 7505; fax: +1 650 246 7776; e-mail: dphillips@portola.com

Current strategies

Current therapeutic strategies for the treatment of arterial thrombosis are based on the well-known receptor systems summarized in Fig. 1. In this simplified diagram, collagen and/or thrombin are designated as the primary platelet agonists. While either agonist is capable of activating platelets, including the activation of the receptor function of GP IIb-IIIa for the binding of fibrinogen and von Willebrand factor (VWF) to initiate platelet aggregation, stable aggregation of platelets is augmented by two autocrine factors generated upon platelet stimulation: ADP, released from platelet dense bodies, and TXA₂, generated by the sequential actions of Cox-1 and thromboxane synthase on the arachidonic acid released from membrane phospholipids. Additional aggregation-dependent secondary mediators include sCD40L and Gas6 plus aggregation-induced tyrosine phosphorylation of GP IIIa and activation of secondary aggregation receptors such as SLAMF1, CD84, Eph kinase and the Gas-6 receptors. Even though the signaling reactions induced by the receptor systems for the platelet stimuli summarized in Fig. 1 are diverse, including those coupled by G_q (PAR-1 and TP), G_i (P2Y₁₂), Syk (GP VI), Shc and talin (GP IIb-IIIa), drugs that target these receptor systems have been designed either to specifically inhibit the receptors themselves [e.g. GP IIb-IIIa antagonists (eptifibatid, abciximab, tirofiban) and P2Y₁₂ inhibitors (clopidogrel, ticlopidine)], to block the generation of the agonists [e.g. the Cox-1 inhibitor aspirin and Factor Xa (FXa) inhibitors [low molecular weight (LMW) heparins]], or to antagonize the agonist itself (e.g. thrombin inhibitors (standard heparin, direct thrombin inhibitors)).

Aspirin

The clinical successes achieved by the current therapies to treat arterial thrombosis have been remarkable. Aspirin was the first

and continues to be the most widely used of these drugs. The trend toward the widespread use of this drug to block arterial thrombosis was first indicated by the findings of the ISIS-2 trial which demonstrated that aspirin reduced mortality from acute MI to a rate that is similar to that of the thrombolytic agent, streptokinase [2]. The data from multiple trials summarized by the Antiplatelets Trialists' Collaboration found a 25% relative risk reduction by aspirin of vascular death, MI or stroke, vs. placebo [3] which led to the widespread adoption of aspirin as standard therapy for primary and secondary prevention of arterial ischemia. This collaboration also reviewed the clinical trials using aspirin to show that low-dose aspirin (75–150 mg daily) is effective for long-term use [4]. While the half-life of aspirin in humans is relatively short (~20 min), its effect persists for the lifetime of an affected platelet in circulation as the drug acetylates Cox-1 at serine-529, located at the active site of the enzyme. Attempts have been made to develop additional drugs that target the thromboxane pathway in platelets including a variety of thromboxane receptor (TP) antagonists, thromboxane A₂ synthase inhibitors, or compounds that combined both functions [5]. Although some of these agents had potent antithrombotic effects in experimental models and preclinical studies, and offered the advantage of inhibiting the TP stimulating activity of prostanoid metabolites in addition to TXA₂ (e.g. isoprostanes, PGH₂), they are not currently used to block arterial ischemia as most were not evaluated in clinically relevant phase III trials [6].

Thienopyridines

The second most widely used of the antiplatelet drugs for chronic use are thienopyridines targeting P2Y₁₂. This class of drugs, which includes clopidogrel, and its predecessor ticlopidine, act via irreversible inhibition of the platelet P2Y₁₂ receptor. Both are prodrugs, requiring hepatic metabolism by




Fig. 1. Current therapeutic strategies for the treatment of arterial thrombosis. The reaction diagram is designed to illustrate the pathways regulated by current therapeutic strategies on unstimulated, discoid platelets (top left), stimulated platelets (top right), and aggregated platelets (bottom). The four antithrombotic drug classes include: (1) Thrombin inhibitors; (2) GP IIb-IIIa antagonists; (3) P2Y₁₂ antagonists; and (4) inhibition of TXA₂ production.

cytochrome P450 isoform 3A4 in order to generate the active metabolite, a transient intermediate which covalently modifies and inactivates the receptor. Ticlopidine has been shown to be efficacious in conditions such as claudication, unstable angina, and cerebrovascular disease [7]. However, the incidence of neutropenia associated with ticlopidine led to the development of a second-generation thienopyridine, clopidogrel, with increased potency and fewer side-effects. In the CAPRIE trial [8], clopidogrel was shown to be more efficacious than aspirin, particularly in high-risk patients (diabetics and those with a history of prior revascularization). Subsequently, the CURE study [9] demonstrated that patients with unstable angina or non-ST segment elevation MI received a 20% relative risk reduction if they were randomized to clopidogrel plus aspirin vs. placebo plus aspirin, and the PCI-CURE substudy [10] showed that this benefit extended to patients undergoing percutaneous intervention (PCI). The slow onset of action of thienopyridines, due to their metabolism requirement, has necessitated the administration of a large loading dose (300 mg) prior to acute procedures, such as PCI, as demonstrated in the CREDO trial, where the maximum benefit of clopidogrel administered with aspirin required a loading dose given at least 6 h prior to the procedure. This study also demonstrated a significant 27% reduction in death, MI and stroke from 1-year administration of clopidogrel plus aspirin following PCI, compared to 1-month dosing [11].

GP IIb-IIIa antagonists

The GP IIb-IIIa antagonists are designed to bind to the integrin on unstimulated platelets and on platelets after stimulation. GP IIb-IIIa is an attractive antiplatelet target as it is (i) on the 'final common pathway' mediating platelet aggregation irrespective of the agonist used to induce platelet activation, (ii) platelet-specific, and (iii) responsible for a variety of aggregation-dependent platelet functions including those in coagulation, inflammation, fibrinolysis and vascular cell proliferation. Three GP IIb-IIIa antagonists have been developed: integrilin, a cyclic heptapeptide modeled after the active site of the disintegrin found in the southeast pigmy rattle snake; abciximab, a Fab fragment of a mouse/human chimeric antibody against GP IIb-IIIa; and tirofiban, a synthetic inhibitor of GP IIb-IIIa. All were designed to be infusible i.v. drugs and are therefore only administered to patients in acute settings who have a high risk of experiencing an ischemic event such as those undergoing PCI (with or without stent placement) or those with symptoms resulting from acute coronary syndrome (ACS) [12]. Use of these drugs has shown a remarkable reduction in death and MI for these indications [13–15].

Anticoagulants

ACS patients, whether undergoing an invasive revascularization procedure or not, are treated with aspirin and antithrombin therapy in the form of unfractionated or LMW heparins. Although unfractionated heparin is effective in reducing clinical

events, a narrow therapeutic index makes it a less than optimal antithrombin for this class of patients. Like their parent anticoagulant, i.e. standard heparin, LMW heparins are indirect antithrombins and utilize antithrombin III to mediate inhibition of thrombin and FXa. As LMW heparins have more predictable pharmacokinetics than standard heparin, they are used in a fixed dose manner. Early trials of the LMW heparin enoxaparin in unstable angina and non-Q wave MI patients demonstrated improved efficacy over standard heparin and the drug has emerged as the most commonly used LMW heparin [16,17]. Fondaparinux, a synthetic pentasaccharide, also utilizes the antithrombin III binding region of heparin and has been found to be an appropriate anticoagulant for prevention of deep vein thrombosis in orthopedic surgery [18]. Unlike enoxaparin, which inhibits both thrombin and FXa, fondaparinux acts only as an indirect FXa inhibitor. Venous thromboembolism prevention trials showed that fondaparinux has a superior efficacy profile to its comparator enoxaparin. Ongoing trials of fondaparinux in ACS patients will show if the concept of attaining superior efficacy by inhibition of FXa alone (vs. the combination of FXa and thrombin) can be achieved in arterial settings.

Combination antithrombotic therapy

Arterial thrombosis developed at sites of spontaneously or mechanically disrupted atherosclerotic plaque is triggered by a multitude of highly thrombogenic materials (i.e. fibrillar collagen and tissue factor). It is the result of complex interrelations between coagulation and platelets orchestrated by local rheological conditions. An emerging strategy in the treatment of arterial thrombosis came with the realization that combinations of antithrombotics provide greater therapeutic benefit than are provided by drugs used singly. Accordingly, the combination aspirin-plus-clopidogrel is rapidly becoming the new standard of care for the management of patients with non-ST segment elevation ACS and in patients undergoing a PCI. In support of this trend, the CURE study demonstrated that aspirin-plus-clopidogrel caused a 20% relative risk reduction of vascular death, MI, and stroke compared with aspirin-plus-placebo [9]. The dual antiplatelet therapy (aspirin-plus-clopidogrel) was also more effective and safer than a combination aspirin-plus-warfarin in coronary artery stenting [19,20]. The remarkable efficacy of the dual anti-platelet therapy has prompted the initiation of several clinical trials in indications as diverse as atrial fibrillation, peripheral arterial disease, peripheral arterial bypass surgery, secondary and high-risk primary prevention, acute ST-segment elevation MI and heart failure [21]. Finally, although anticoagulants were routinely used in the development of antiplatelet agents, analysis of these data shows that these combinations often provided a clinical benefit that was greater than anticipated. We and others have used thrombosis models to evaluate synergisms between various pathways. Because TXA₂ and ADP activate different pathways, it was anticipated that combinations of inhibitors of the two pathways would confer a




Fig. 2. Thrombosis defect in platelets from the diYF mouse. Unanticoagulated blood from a control mouse (WT) or from a mouse harboring the diYF mutation in GP IIIa (Y747F, Y759F) was perfused for 2.5 min through a chamber coated with type III collagen at 871 s^{-1} and stained. The view is en face.

greater protection against thrombotic events. However, rather than an additive effect, the two drugs used together were synergistic [22,23]. A potent synergism between clopidogrel and anticoagulants [using either a direct thrombin inhibitor (Bivalirudin), a FXa inhibitor or a synthetic LMW heparin] has also been observed (Fig. 2) [23,24]. While these results may have a simple explanation, resulting from the inhibition of fibrin formation by the anticoagulant, where fibrin contributes to thrombus stability under arterial shear rate [25], the emerging data indicate that the anti-thrombotic synergisms may originate from the complementation of the signaling pathways in platelets. Activation of the receptor function of GP IIb-IIIa is optimal when engagement of $G_{12/13}$ or G_q signaling pathways is combined with G_i stimulation [26,27]. This combination occurs when either TXA_2 (TP) receptor ($G_{12/13}$, G_q), protease-activated receptor-1 (PAR-1) ($G_{12/13}$, G_q), and also potentially P2Y_1 (G_q) [28] is allowed to synergize with that of P2Y_{12} (G_i). Note that this model might also explain the synergism between aspirin and clopidogrel. A different explanation may stem from the inhibition of the different PI-3 kinase enzymes present in platelets. For example, P2Y_{12} engagement by ADP stimulates PI-3 kinase- γ whereas the engagement of either $\text{Fc}\gamma\text{RIIA}/\text{GPVI}$, PAR-1, TP, or GP IIb-IIIa lead to PI-3 kinase α or β -activation. This could indicate that any combination therapy that would block PI-3 kinase α , β or γ would confer a strong antithrombotic efficacy.

Limitations of current antithrombotics

Even with the remarkable successes that have been achieved with currently available antithrombotics in the prevention of arterial thrombosis, limitations of this class of drugs do exist. It is valuable, therefore, to consider additional strategies currently available to design new drugs that address these limitations. That there is room for improvement is readily apparent from analysis of current trials. For example in SYNERGY, a trial with more than 10 000 high risk non-ST segment elevation ACS patients treated with heparin or LMW heparin, aspirin, and, as determined by the physician, clopidogrel and/or GP IIb-IIIa antagonists, approximately 15% of all patients still

experienced death or non-fatal MI within 30 days of treatment [29]. Except for aspirin, perhaps the biggest limitation of these drugs is that the dose used is less than optimal for the treatment of thrombosis. The drugs were typically titrated in preclinical development studies to arrive at a dose that had a significant inhibition of thrombosis without undue bleeding. While higher doses of the drugs had better antithrombotic activity, this always created a bleeding risk. Although the dose selected required some refinement in subsequent clinical studies, this same strategy was employed.

Aspirin non-responsiveness

While aspirin is used at a dose (e.g. $70\text{--}325 \text{ mg day}^{-1}$) that yields near 100% acetylation of Cox-1 in most individuals, it has been recognized for several years that individuals treated with aspirin still experience thrombotic events. The interpretation of this observation has been controversial. One could argue that aspirin is a comparatively 'weak' inhibitor in that it only blocks the production of TXA_2 , an autocrine factor that supplements the activities of the primary platelet agonists. Indeed, aggregation reactions using platelets from individuals taking aspirin appear normal when high concentrations of primary agonists are used. However, it has become clear from recent pharmacodynamic and biochemical studies that platelet responses normally blocked by inhibition of Cox-1 (the target of aspirin in platelets) are still present in some individuals, an observation that has led to the concept of 'aspirin resistance' or 'aspirin non-responders' [30–32]. Studies of platelet aggregation in aspirin-treated CVD patients, both by traditional ADP and arachidonic acid induced aggregation studies and by platelet function analyzers such as PFA100, show that 5–10% of the individuals can be classified as aspirin non-responders [31]. Additional studies demonstrated a threefold higher risk of major cardiovascular adverse events in the same patient population [30]. As the incidence of aspirin non-responders had been shown to be higher in patients who undergo coronary artery bypass graft (CABG), a recent study evaluated the functional and biochemical responses to aspirin on subsequent days following a CABG procedure in a small number of patients [33]. The study demonstrated that platelets from these patients after CABG did not completely respond to aspirin *in vitro*, and that while Cox-1 levels in platelets remained constant 10 days following the procedure, there was a pronounced increase in the level of Cox-2, which is ~ 170 -fold less sensitive to aspirin inhibition, especially at 5 days post-procedure. This may be reflective of the increased platelet turnover following cardiopulmonary bypass, and the increased level of Cox-2 could generate critical amounts of TXA_2 , in spite of aspirin treatment, providing a possible explanation for aspirin non-responsiveness. Other mechanisms proposed as contributing factors to aspirin non-responsiveness include use of NSAIDs, which block acetylation by aspirin [34] and polymorphisms of platelet genes (Cox-1 or GP IIb-IIIa) [35], and non-compliance. However, the answer may be totally unexpected such as the activation of a deacetylase. Clearly, further

studies are needed to better define the underlying mechanisms of this phenomenon. Aspirin use is also contraindicated in a significant population of patients, i.e. those with gastrointestinal bleeding and those with aspirin-induced asthma.

Clopidogrel non-responsiveness

The primary limitation of clopidogrel is that this drug demonstrates weak and somewhat variable inhibition of P2Y₁₂ [36]. Following a 600-mg loading dose of clopidogrel, the extent of inhibition of ADP-induced aggregation (5 μ M ADP) varied from 33% to 78% in healthy individuals, at 6 h post-dosing [37]. This effect is further exaggerated in patients undergoing PCI/stent placement [38,39]. The antithrombotic effect of clopidogrel is likely to be dependent on a number of factors including but not limited to variations in P450s, polymorphisms of the P2Y₁₂ receptor and receptor signaling pathways. Measurements of platelet aggregation and markers of platelet activation (GP IIb-IIIa and P-selectin detection by specific antibodies) show that clopidogrel resistance is detected in 31% of the patients on day 5 and 15% of the patients on day 30 of the treatment regimen [40]. A prospective study of PCI patients with non-ST segment elevation MI showed that up to 25% of the patients were resistant to clopidogrel [41]. When the patients were stratified into quartiles based on resistance to ADP-induced platelet aggregation, the most resistant patients had a 40% adverse event rate during a 6-month follow-up period so they are obviously being denied adequate protection based on inhibition of P2Y₁₂. It has also been reported that the antiplatelet activity of clopidogrel is blocked in patients treated with a widely used cholesterol lowering medication (atorvastatin) which is undoubtedly linked to the metabolism requirement for efficacy [42]. A second limitation of these drugs stems from their irreversible mechanism of action, which inactivates the P2Y₁₂ receptor for the lifetime of the platelet. While this is not a particular problem with aspirin, which is also irreversible, this feature has led to limited use of clopidogrel before PCI in patients who are at increased risk of undergoing CABG procedures, as the risk for bleeding following clopidogrel treatment requires postponement of the procedure for 5–7 days, or transfusion of large numbers of platelets during the procedure.

Limitations of anticoagulants – i.v.

Each anticoagulant has evolved unique issues. Replacement of unfractionated heparin has had mixed success. Although the current ACC/AHA guidelines (2002) prefer enoxaparin over unfractionated heparin, recent data do not support this preference. In the SYNERGY trial, enoxaparin was found not to be superior to unfractionated heparin [29]. In A to Z, for patients on a GP IIb-IIIa antagonist (tirofiban) and aspirin, enoxaparin was found to be non-inferior to standard heparin [43]. LMW heparins also have a narrow safety window. For example, in unstable angina patients treated with enoxaparin, a 25% dose increase in therapeutic level (1.25 mg kg⁻¹ vs.

1 mg kg⁻¹) produces an unacceptable number of bleeding events [44]. Direct thrombin inhibitors such as angiomas have also been studied as replacements for unfractionated heparin. In the REPLACE 2 trial of PCI, angiomas and provisional GP IIb-IIIa inhibitor compared favorably to heparin plus GP IIb-IIIa [45]. The primary end point of the trial combined efficacy and safety parameters and the angiomas arm of the trial was statistically not inferior to the heparin arm. However, the benefit related to reduction of bleeding with the use of angiomas is questionable. In REPLACE 2, the control group was likely to have been over anticoagulated as the observed ACT values in the heparin arm were higher than the recommended ACT range (200–300 s) for use of heparin in conjunction with GP IIb-IIIa. As angiomas is substantially more expensive than standard heparin, economic considerations also contribute to its limited use in PCI patients.

Limitations of anticoagulants – oral

Warfarin is the only anticoagulant in chronic use. While the drug provides tremendous benefit to affected individuals, its anticoagulant response is influenced by a variety of factors such that >50% of patients are usually outside of the therapeutic range. Due to the large variability in the anticoagulant effect of warfarin and its narrow therapeutic index, a large unmet clinical need exists for an anticoagulant with predictable fixed-dose usage. The need for a warfarin substitute has led to numerous drug development projects that have focused on inhibitors of coagulation proteases that specifically inactivate the protease active site. Ximelagatran, an oral thrombin inhibitor, was the first to show that the strategy of direct coagulation protease inhibition does translate into effective anticoagulation and leads to antithrombotic activity in deep vein thrombosis and atrial fibrillation patients [46]. Two studies suggest that ximelagatran is at least as effective as warfarin in preventing stroke in high-risk patients with atrial fibrillation [47]. The studies also showed that there are incidences of increase in liver enzymes which would require surveillance for potential liver toxicity in future patients. Unfortunately, safety problems of ximelagatran related to serious liver toxicity has led the FDA to recommend against approval of this thrombin inhibitor. While several new experimental agents with the potential to be an effective and low variability anticoagulant have been evaluated in clinical trials, none of these are available for therapeutic use, so the search for a warfarin replacement remains a work in progress.

PAR-1

While drugs that inhibit thrombin or prevent its formation are a mainstay in the armamentarium used for the treatment of arterial thrombosis, as of this writing, no efficacy trials have been performed to determine whether antagonists of PAR-1, the thrombin receptor on platelets, could provide a therapeutic benefit. Potent and selective PAR-1 antagonists capable of inhibition of thrombin-induced platelet aggregation have been

reported in the literature. Peptide mimetic antagonists such as RWJ-58259 are effective in models of thrombosis and vascular injury and could have potential as therapies for treating thrombosis and restenosis [48]. An oral PAR-1 antagonist, E-5555, is being developed as a drug candidate for ACS but definitive clinical efficacy trials have not been reported [49].

Future directions for antithrombotic drug development

New antithrombotics are required not only to overcome the limitations of the current drugs to better manage arterial ischemia, but also to address the inflammatory activities of platelets which contribute to progression of atherosclerotic disease. The successes and limitations of current therapies coupled with the advances made in our understanding of platelet biology are instructive in the design of new drugs to more effectively regulate validated targets, in the identification of new targets that may safely provide increased benefit and in the development of the proper combination of antithrombotics for the various arterial ischemic indications.

Agonist receptors

While platelets are activated by numerous agonists acting on multiple receptors, the only validated agonist receptor for drug discovery is P2Y₁₂. The requirements for improvements over clopidogrel are clear – more potent inhibition of P2Y₁₂; less variability of inhibition between different patients; no requirement for metabolism resulting in less delay in onset to action; and quicker recovery of platelet function following discontinuation of use. While these requirements can most likely be best achieved by an orally available reversible P2Y₁₂ antagonist, preclinical data indicate three promising candidates in development with different properties. One is cangrelor (AR-C69931MX), a nucleotide, intravenous compound that reversibly antagonizes P2Y₁₂ [50]. AZD-6140, an orally available direct-acting P2Y₁₂ antagonist [51], is presently being evaluated in phase II clinical trials. Prasugrel (CS-747), a thienopyridine prodrug similar to clopidogrel which is more rapidly converted to the active metabolite than is clopidogrel, has completed phase II trials and will be evaluated in phase III trials in ACS patients [52].

ADP also acts on P2Y₁, a G_q coupled receptor. Studies using either selective antagonists of P2Y₁ or P2Y₁₂, as well as gene-targeting strategies [53–56] have demonstrated distinct roles for these two ADP receptors. P2Y₁ is responsible for initiation of aggregation to ADP [55], while P2Y₁₂ is critical for amplification of the aggregation response by released ADP, and for stabilization of platelet aggregates and the growing thrombus [57]. In addition to the different roles of these two receptors in initiation and stabilization of thrombus growth, one could argue that the selective tissue distribution of P2Y₁₂ (platelets, megakaryocytes and glial cells), vs. P2Y₁ (which is ubiquitously expressed) makes it the preferred drug target. Although there are selective P2Y₁ antagonists which have been used as *in vitro* tools [58,59], none of these have been clinically evaluated as yet,

and may not have suitable pharmacokinetic properties to be viable drug candidates.

Secondary aggregation receptors

While the initial interaction of platelets during thrombosis is dependent upon GP IIb-IIIa, it has become apparent that signaling reactions initiated by platelet–platelet contact are required for thrombus stability. Several mediators of aggregation-induced signals have been identified. One is GP IIb-IIIa itself which becomes tyrosine phosphorylated and also associates with numerous signaling and cytoskeletal proteins following platelet aggregation. The importance of the ‘outside-in’ signaling in the enhancement of platelet aggregation was demonstrated by the generation of knock-in mice where tyrosine residues Y747 and Y759 were mutated to phenylalanine [60]. The so-called DiYF mice displayed selective impairment of outside-in signaling resulting in the formation of unstable aggregates. In addition, as shown in Fig. 2, *ex vivo* perfusion chamber experiments on type III collagen have shown that the DiYF mouse has defective thrombosis. Another protein involved in secondary platelet aggregation is CD40L, a tumor necrosis factor family member mainly expressed on activated T cells and platelets [see 61]. CD40L is cryptic in unstimulated platelets, but rapidly becomes exposed on the platelet surface after stimulation where it is subsequently cleaved, producing a soluble hydrolytic product termed sCD40L [61]. We have shown that mice lacking CD40L have a thrombosis phenotype and that normal thrombosis is regained upon infusion of sCD40L [62]. Interestingly, sCD40L, in addition to being a ligand for CD40, is also a ligand for GP IIb-IIIa, a reaction that depends upon its KGD sequence, a known GP IIb-IIIa binding motif. sCD40L also triggers outside-in signaling by tyrosine phosphorylation of GP IIIa, a reaction which is defective in the platelets from the DiYF mouse [63]. While inhibition of primary platelet aggregation and this secondary aggregation mechanism are both inhibited by GP IIb-IIIa antagonists, a potential drug target is the metalloproteinase responsible for CD40L cleavage. Another protein released upon platelet activation that functions to consolidate platelet thrombi is Gas6. Gene targeting of Gas6 also demonstrates a thrombosis phenotype [64]. Gas6 binds to three receptors on platelets, Tyro3, Axl and Mer, but genetic targeting of any one unexpectedly inhibits Gas6-induced platelet stimulation. However, as Gas6 also induces tyrosine phosphorylation of GP IIIa, apparently by a mechanism independent of binding to the integrin, it has been proposed that Gas-6 signaling could be therapeutically regulated through inhibition of Gas6-GP IIb-IIIa cross-talk [65].

Platelet–platelet contacts induce the activation of additional signaling mechanisms which are involved in aggregate stability. One involves Eph kinases and ephrins, specifically EphA4 and ephrinB1, which through receptor ligand interactions on the platelet surface enhance the binding of GP IIb-IIIa to immobilized fibrinogen in the presence of physiological agonists [66]. Recent work from our laboratory using both

oligonucleotide-based microarray analyses and mass spectrometric proteomics techniques has identified two additional receptor families that are involved. One involves two members of the SLAM family of adhesion receptors, SLAM and CD84; the other involves a novel protein termed PEAR1 (N. Nanda, M. Hart and D.R. Phillips, unpublished data). All proteins are exposed on the surfaces of unstimulated platelets and signal secondary to GP IIb-IIIa-mediated platelet-platelet contacts by becoming tyrosine phosphorylated. Therapeutic targeting of one or more of these secondary aggregation receptor systems in platelets is an attractive possibility as they appear to have a greater effect on thrombosis than they do on hemostasis.

Adhesion receptors

Several platelet adhesion receptors have been identified but we will focus on GPVI and GPIIb α , the adhesion receptors that are not only involved in the adhesion of platelets to the highly thrombogenic fibrillar collagens in the vessel wall exposed following vascular injury but also contribute to platelet activation. Both receptors are attractive drug discovery targets as both are platelet specific. Under the high shear rates encountered in coronary and carotid arteries, the binding of VWF to the collagen surface triggers a transient interaction with GPIIb α that allows for a more stable interaction of the platelet with the collagen surface via at least two collagen receptors, integrin $\alpha_2\beta_1$ and GPVI. Recent findings indicate that GPIIb α and $\alpha_2\beta_1$ preferentially contribute to the adhesion process whereas engagement of GPVI triggers signaling events leading to platelet activation [67]. GPVI is non-covalently associated with the Fc receptor γ -chain and signals through the platelet via stimulation of multiple non-receptor tyrosine kinases. Interestingly, part of this signaling pathway may be common to GPIIb α activation, and signals coming from these two receptors contribute to the activation of the receptor function of GP IIb-IIIa and platelet aggregation. Modulation of the GPVI receptor function is becoming an attractive target as platelets from GPVI-deficient animals, human platelets expressing low levels of GPVI, or platelets treated with a GPVI antibody, while unable to support thrombus growth [67–69], are nonetheless able to adhere on the collagen surface (via $\alpha_2\beta_1$ integrin and potentially another collagen receptor [70]) minimizing the effects on hemostasis [71,72]. GPIIb α is a high shear rate-dependent thrombosis receptor that affects recruitment of platelets at sites of vascular injury (on the collagen present in the subendothelium and on adhering platelets) with minor impact on venous thrombotic process. Modulation of the VWF/GPIIb α axis has been the subject of many investigations with promising animal experimental results, but severe thrombocytopenia has been associated with the use of antibodies against GPIIb α , thus reducing the general interest of the scientific community for several years. Nevertheless, novel strategies targeting the VWF/GPIIb α axis through snake venom proteins cleaving GPIIb α , VWF peptides or antibodies against VWF are reviving this strategy.

Signaling pathways

The extensive repertoire of platelet functions, while initiated by receptors, is regulated by signal transduction pathways. While these pathways have not been known as drug discovery targets, two observations suggest that they are worthy of consideration. First, the remarkable success of Gleevec, an Abelson tyrosine kinase inhibitor, has proven efficacy in the treatment of chronic myelogenous leukemia and other cancers. While the multiple functions of this kinase in diverse cell types predicted toxicity, clinical data have shown that the benefits far outweigh the liabilities. The success of this drug suggests that signal transduction pathways, though redundant for multiple signaling systems in diverse cell types, are worthy of consideration as therapeutic drug discovery targets. This conclusion is supported by the analysis of numerous gene-targeted mouse strains which have led to the surprising conclusion that the phenotype achieved by the disruption of any specific gene is often limited, even for genes involved in signal transduction. A second observation is that the platelet stimuli often induce diverse responses. For example, any one of the primary platelet agonists are capable of producing a spectrum of responses which could have pathological implications, e.g. aggregation caused by the activation of the receptor function of GP IIb-IIIa, expression of procoagulant activity of prothrombinase or FXase to catalyze the production of thrombin, generation of vasoactive substances such as TXA₂ and serotonin to induce vasoconstriction, release of proinflammatory proteins like sCD40L, RANTES and TGF- β to affect vascular inflammation including the progression of atherosclerosis, the release of growth factors such as PDGF to affect vascular remodeling, and the activation of secondary aggregation receptors such as SLAM, CD84 and the ephrins to stabilize thrombi and cause vascular occlusions. As many of these responses would be expected to be regulated by a specific pathway, it is reasonable to expect that these responses could be individually regulated. If true, this approach could inhibit platelet-dependent pathologies without compromising primary hemostasis. One potential example is PI-3 kinase and the regulation of the adhesive function of GP IIb-IIIa [73].

The roles of secondary signaling events downstream of platelet surface receptors have been elucidated through gene-targeting studies in mice, and subsequent evaluation of their platelet phenotypes using both *in vitro* and *in vivo* techniques. As certain key platelet agonists such as ADP, thrombin, and TXA₂ all activate platelets through G protein-coupled receptors, genetic targeting of individual α -subunits of G proteins has been a successful strategy in studying platelet signaling downstream of receptor activation. Characterization of platelets from mice lacking G_q, G₁₂ and G₁₃, identified these three proteins as key mediators for ADP and TXA₂ receptors [27,74–76]. Targeting of additional subunits (G₂, G₁₃, and G₁₂) showed little or no effect on platelet phenotype, which could be due to lack of coupling of these subunits to critical platelet receptors, or due to redundancy in the signaling pathways. Thrombin signaling has been shown to be affected by lack of G_i, G_q, and

G₁₃, to varying extents. In mice lacking both G_q and G₁₃, no platelet activation was possible by ADP, TXA₂, or thrombin [27], suggesting that at least G_q or G₁₃ is required to induce some activation, and that activation of G_i-type proteins alone is not sufficient for activation of mouse platelets. In these G_q/G₁₃ double-deficient platelets, adhesion of platelets to collagen was not affected; however, aggregation in response to collagen fibrils as well as formation of stable aggregates on collagen-coated surfaces was completely eliminated. In addition to targeting of G protein subunits, targeting molecules involved in kinase signaling pathways have resulted in mice with impaired platelet functions, which is not unexpected given the fact that kinases of the src family (Csk, src) have been shown to be physically associated with the cytoplasmic domain of GP IIb-IIIa [77]. The platelet phenotype of mice lacking the tyrosine kinase syk, critical for downstream signaling through the collagen receptor GPVI, and also activated during outside-in signaling and activation of GP IIb-IIIa, exhibited defects in platelet activation induced by ADP ± epinephrine [78]. Mice lacking the adapter protein SLP-76, which is on the syk signaling pathway, have been shown to have defects in GP IIb-IIIa signaling and collagen receptor responses [79]. The important role of downstream signaling molecules identified through gene-targeting studies may provide new opportunities for therapeutic intervention for blockade of platelet activation, thrombus formation, and adhesion.

Evolving paradigm on the relationship of thrombosis to inflammation

Not only are platelets critical players in mediating thrombosis, but recently their role in inflammation has become more appreciated. Although it is widely accepted that inflammatory activities that orchestrate the progression of atherosclerosis are derived from 'traditional' inflammatory cells such as monocytes and neutrophils, emerging data suggest that products released from platelets during thrombosis are actively involved in this process and that platelets are a primary source of inflammatory proteins within the circulation. For example, Schober et al. [80] reported that platelets deposit RANTES onto endothelial cells in the injured vessel wall, and that this interaction is mediated by P-selectin, a surface receptor mediating the attachment of platelets to leukocytes and endothelium. Additional work from this laboratory showed that infusion of activated platelets into the apoE^{-/-} mouse greatly enhanced the rate of atherosclerotic lesion progression [81]. The exposure of P-selectin following platelet activation is a key mediator of platelet-leukocyte interaction, and facilitates atherosclerotic lesion development, as demonstrated by Burger and Wagner [82]. In patients with acute MI, platelet-leukocyte interaction is increased compared with controls, and P-selectin levels have been shown to remain increased for at least a month following initial presentation in ACS patients with non-ST segment elevation [83], and even in patients with stable coronary artery disease. Antiplatelet therapy with a P2Y₁₂ antagonist plus aspirin was shown to decrease platelet-monocyte interactions that occur after

coronary stenting [84], an effect not observed with anticoagulants plus aspirin, suggesting that P2Y₁₂ antagonism had an anti-inflammatory effect distinct from its antithrombotic mode of action. Targeting of CD40L, another platelet-derived inflammatory protein, either through a blocking antibody [85] or via gene-targeting [86], greatly inhibited lesion progression in either LDLR^{-/-} or apoE^{-/-} mice, respectively. Soluble CD40L (sCD40L), the shed hydrolytic product of CD40L, 95% of which is platelet-derived, has been shown to be a primary risk factor for atherosclerosis/thrombosis [87]. In addition, the binding of platelet-derived sCD40L to endothelial cells can lead to the expression of tissue factor, a potent procoagulant. Clopidogrel has been shown to inhibit ADP-induced CD40L expression, and to lower CD40L levels in patients undergoing PCI [88], while aspirin does not. Thus, some forms of antiplatelet therapy, including P2Y₁₂ inhibition, can inhibit platelet pro-inflammatory responses. Finally, recent data show that targeting of GP Ib-IX-V complex, a platelet adhesion receptor, in the apoE^{-/-} mouse blocks leukocyte recruitment and the development of atherosclerotic lesions [89]. Thus, studies of the role of platelets in inflammation may provide new potential targets for CVD through inhibition of atherosclerosis and/or thrombosis.

Platelet monitoring

The pharmaceutical industry and drug approval agencies expect that the early inroads into personalized medicine in the administration of selected chemotherapies will ultimately extend to all drug classes. These early examples include screening for HER2 positive individuals in the treatment of breast cancer with Herceptin and the screening for EGF receptor for the treatment of lung cancer with gefitinib [90]. More generally, by the time this manuscript is published, the CYP450 screen will most likely be available to detect the various isoforms of P450 which will be useful in projecting drug levels in individuals treated with a wide variety of drugs. While such strategies will be of value in selecting and dosing antithrombotics, particularly for chronic use, access to blood of patients treated with antithrombotics continues to provide the best opportunity for monitoring the effect of any therapy or any combination of therapies on each patient being treated. While suitable assays are available to monitor anticoagulants, the monitoring of antiplatelet drugs is not routinely performed.

Evaluations of data from the current antithrombotics and their limitations have defined the assays required to bring personalized medicine to the patient treated with antiplatelet drugs to prevent arterial thrombosis. First, platelet function should be monitored in the context of thrombosis. Platelet thrombosis *in vivo* is initiated by adhesive proteins exposed on the vessel wall and stable thrombi result following adhesion, activation, aggregation, and thrombus stabilization, all occurring under conditions of shear. As thrombus stability is one of the issues, continuous monitoring of thrombus formation is essential for determining the effects of drugs that effect targets

involved in thrombus stability, for example, in prostanoid metabolism and in ADP release. Although methods currently available such as light transmittance aggregometry, the Ultegra Rapid Platelet Function Assay and platelet activation markers such as P-selectin expression are effective in monitoring the ability of end products of any one of these pathways to activate platelets, they are ineffective in monitoring thrombosis, the physiological response of platelets to thrombogenic surfaces under shear. Furthermore, while the PFA-100 device is capable of monitoring the time required for a platelet plug to form in apertures coated with ADP or collagen, it does not provide a continuous monitoring of the thrombotic process. A second requirement for the personalized monitoring of drugs to prevent arterial thrombosis is that it should not only be responsive to diverse drug classes but also be capable of determining the net effect on thrombosis achieved by combinations of antithrombotic therapies. At the present time, the four drug classes used to treat patients at risk for arterial thrombosis, aspirin, GP IIb-IIIa antagonists, clopidogrel and anticoagulants, are used in combinations not properly evaluated for their net effect on thrombosis. Drugs against additional platelet and coagulation protein targets will become available. Required are technologies that will readily permit evaluation of how combinations of these drugs affect thrombosis in patients receiving these drugs. A third requirement is that the method must be capable of monitoring individual differences in response to therapy. As outlined above, individual differences in response to aspirin and clopidogrel have been observed – differences which appear to affect clinical outcome. While currently available techniques such as light transmittance aggregometry, Ultegra, or PFA-100 are useful in monitoring responsiveness to both of these drugs as monotherapy, they are ill-suited to measure individual differences when combinations of drugs are employed, the emerging norm. It is also anticipated that this problem would be amplified when drugs against additional targets would be introduced. Finally, as the anticoagulants routinely used in blood collection such as citrate or direct thrombin inhibitors such as PPACK affect thrombosis, the monitoring method must be capable of assaying non-anticoagulated samples of blood. Perhaps the best example of anticoagulant interference in antithrombotic monitoring is in the development of GP IIb-IIIa antagonists where citrate anticoagulation markedly overestimates antiaggregatory activity [91]. The optimal method for monitoring individual thrombotic potential must be capable of either performing the assay in the absence of anticoagulants or of being able to determine how any given anticoagulant affects the assay.

Use of perfusion chamber technology has perhaps provided the best hope of measurement of the thrombotic potential of individuals being treated with combinations of antithrombotic drugs. Perfusion chambers were designed 30 years ago in order to characterize the thrombotic process under shear conditions. The different types of perfusion chambers described in the literature can be classified according to their geometry (circular, annular, flat chambers) or the surfaces (blood vessels, isolated proteins) exposed to flowing blood. These techniques confer the

advantage of studying platelet interactions with a thrombogenic surface under specific conditions of shear rate with either non-anticoagulated or anticoagulated blood. Major contributions to the field of thrombosis have originated from use of perfusion chambers. For example, the critical role of VWF and its interactions with GP Ib and GP IIb-IIIa to mediate platelet adhesion and thrombus growth under arterial shear rates, the involvement of GP VI and of the integrin $\alpha 2\beta 1$ in mediating platelet adhesion and activation on collagen. Inhibitors of P2Y₁₂ and Cox-1 have also demonstrated antithrombotic activities in this system [92]. However, perfusion chambers are mostly utilized by academic institutions or by pharmaceutical and biotechnology companies in order to identify or validate targets and to develop antithrombotic drugs. Several limiting factors have prevented their use as a bedside device for monitoring drug efficacy in clinical trials – the skill required to determine thrombus size was not readily available in clinical settings; quick readouts for the patient were not available; end point quantifications left investigators without knowledge of the kinetics of thrombus formation, the more critical information.

Results from several laboratories, however, have made progress in modifying these devices to circumvent these difficulties. Figures 3 and 4 illustrate the utility of monitoring the kinetics of thrombosis in perfusion chamber assays. In one instance, using non-anticoagulated samples of blood, we have shown that inhibition of P2Y₁₂, Cox-1, or FXa did not significantly reduce thrombus size after a 4-min perfusion period over a collagen-coated surface. However, when more than one of these targets was inhibited, pronounced anti-thrombotic activity was observed. In another experiment, when human blood was anticoagulated with an FXa inhibitor and perfused through a chamber in the real-time assay, we observed that P2Y₁₂ antagonism with clopidogrel did not affect the




Fig. 3. Synergism between P2Y₁₂ antagonism, Factor Xa inhibition and aspirin. As indicated unanticoagulated blood was treated with an inhibitor for P2Y₁₂ (100 µM 2MeSAMP) or Factor Xa (10 µM C921–78). Aspirin-treated was from aspirin-treated individuals. The treated blood was perfused through a chamber coated with type III collagen at 1000 s⁻¹ for 4 min and quantified as described [23].




Fig. 4. Synergism between aspirin and P2Y₁₂ inhibition in blocking thrombus growth. Blood from a control individual, an individual treated with clopidogrel, or an individual treated with aspirin and clopidogrel was anticoagulated with a Factor Xa inhibitor, treated with rhodamine 6G to label platelets, and then perfused through a chamber coated with type III collagen at 1000 s⁻¹. The continuous accumulation of fluorescence was used to quantify platelet thrombus formation.

initial thrombus growth triggered by fibrillar type III collagen under arterial shear rates. However, clopidogrel caused the thrombi formed during the first 3 min of perfusion to dissociate. Control thrombi formed in the absence of clopidogrel were stable and continued to grow. This demonstrates the limitations of end point analysis as the measured anti-thrombotic activity is dependent on the time of analysis. Retrospectively, this explains an apparent discrepancy found in the evaluation of the phenotype of P2Y₁₂^{-/-} mice [57]. P2Y₁₂^{-/-} mice demonstrate a cyclic thrombotic process *in vivo*, but only a qualitative difference (i.e. more loosely packed thrombi) was observed after perfusion of non-anticoagulated blood for 2.5 min over type III collagen. Evaluation of mono- and combination therapies in this assay confirmed the antithrombotic efficacy of the different anti-platelet therapies, with GP IIb-IIIa antagonists being inhibitors of thrombus growth, aspirin and P2Y₁₂ antagonist destabilization agents, the combination aspirin + P2Y₁₂ antagonism showing a faster destabilization activity (Fig. 4). Thus, perfusion chamber technology is suited to meet the requirements of personalized medicine for individuals receiving antithrombotic therapies as the measurement is on thrombosis, it is responsive to diverse drug classes and it can be performed in the absence of anticoagulants. Future discoveries are required to adapt such technologies to devices that are readily available to individual patients. Finally, several laboratories have reported measurements of the inflammatory activities of platelets, e.g. sCD40L plasma levels, P-selectin expression, formation of platelet-leukocyte complexes. As recent data show that the inflammatory activity of platelets is important in the progression of atherosclerosis, it would also be desirable to develop methods to rapidly quantitate the platelet inflammatory activity in patients.

Application of this personalized medicine approach to antithrombotic therapies does have significant hurdles to overcome before it can be used to reliably modify therapy.

First, a bedside monitor of the thrombotic potential of individual patients needs to be developed. Second, recognizing that individuals will undoubtedly be heterogeneous with respect to vessel wall thrombogenicity, including the local shear environment, results using this device need to be correlated with clinical outcomes.

Conclusions

The current repertoire of drugs for the treatment of patients at risk for arterial thrombosis (e.g. ACS, diabetes, poststroke, peripheral artery disease, post-AMI) currently includes four classes of drugs – aspirin, GP IIb-IIIa antagonists, thienopyridines, and anticoagulants. Although each of these drug classes has proven efficacies for different indications, each has limitations that continue to permit thrombotic events during their use. In addition, emerging data suggest that a significant percentage of individuals treated with aspirin or clopidogrel do not receive the expected therapeutic benefit from therapy because of a decreased responsiveness by their platelets. Future directions in addressing these limitations will proceed in two parallel directions. On the one hand, it can be anticipated that new drugs, either offering improvements against known, validated targets, or against recently identified targets, will be forthcoming. Recognizing that platelets are now known to be directly involved in vascular inflammation including that which leads to the progression of atherosclerotic disease, it can be anticipated that some of these new therapeutic strategies will not only better address arterial thrombosis, but also inhibit the ability of platelets to deliver inflammatory proteins and growth factors which affect atherosclerotic lesion development. On the other hand, it has now become apparent that improvements are required in the devices used to monitor the thrombotic potential of individuals receiving therapy, both for the development of new antithrombotic drugs and to measure the effectiveness of combined antithrombotic therapies. It would appear that the most effective device is that which measures thrombosis in real time, is accessible to the patient at the point of drug administration, and can be performed in the absence of anticoagulation. Such a device would be capable of monitoring the activities of new classes of antithrombotics, of measuring variances of individual responses, and in evaluating the effectiveness of combined antithrombotic therapies.

References

- 1 Fox CS, Coady S, Sorlie PD, Levy D, Meigs JB, D'Agostino RB Sr, Wilson PW, Savage PJ. Trends in cardiovascular complications of diabetes. *J Am Med Assoc* 2004; **292**: 2495–9.
- 2 Fox CS, Coady S, Sorlie PD, Levy D, Meigs JB, D'Agostino RB Sr, Wilson PW, Savage PJ. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. *Lancet* 1988; **2**: 349–60.
- 3 Fox CS, Coady S, Sorlie PD, Levy D, Meigs JB, D'Agostino RB Sr, Wilson PW, Savage PJ. Collaborative overview of randomised trials of antiplatelet therapy – I: prevention of death, myocardial infarction,

- and stroke by prolonged antiplatelet therapy in various categories of patients. Antiplatelet Trialists' Collaboration. *Br Med J* 1994; **308**: 81–106.
- 4 Fox CS, Coady S, Sorlie PD, Levy D, Meigs JB, D'Agostino RB Sr, Wilson PW, Savage PJ. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *Br Med J* 2002; **324**: 71–86.
 - 5 Dogne J-M, de Leval X, Delarge J, Masereel B. Recent developments of thromboxane modulators. *Expert Opin Ther Patents* 2001; **11**: 1663–75.
 - 6 Jneid H, Bhatt DL. Advances in antiplatelet therapy. *Expert Opin Emerg Drugs* 2003; **8**: 349–63.
 - 7 Janzon L. The STIMS trial: the ticlopidine experience and its clinical applications. Swedish Ticlopidine Multicenter Study. *Vasc Med* 1996; **1**: 141–3.
 - 8 Janzon L. A randomised, blinded, trial of clopidogrel versus aspirin in patients at risk of ischaemic events (CAPRIE). CAPRIE Steering Committee. *Lancet* 1996; **348**: 1329–39.
 - 9 Yusuf S, Zhao F, Mehta SR, Chrolavicius S, Tognoni G, Fox KK. Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. *N Engl J Med* 2001; **345**: 494–502.
 - 10 Mehta SR, Yusuf S, Peters RJ, Bertrand ME, Lewis BS, Natarajan MK, Malmborg K, Rupprecht H, Zhao F, Chrolavicius S, Copland I, Fox KA. Effects of pretreatment with clopidogrel and aspirin followed by long-term therapy in patients undergoing percutaneous coronary intervention: the PCI-CURE study. *Lancet* 2001; **358**: 527–33.
 - 11 Steinhubl SR, Berger PB, Mann JT III, Fry ET, DeLago A, Wilmer C, Topol EJ. Early and sustained dual oral antiplatelet therapy following percutaneous coronary intervention: a randomized controlled trial. *J Am Med Assoc* 2002; **288**: 2411–20.
 - 12 Scarborough RM, Kleiman NS, Phillips DR. Platelet glycoprotein IIb/IIIa antagonists. What are the relevant issues concerning their pharmacology and clinical use? *Circulation* 1999; **100**: 437–44.
 - 13 O'Shea JC, Madan M, Cantor WJ, Pacchiana CM, Greenberg S, Joseph DM, Kitt MM, Lorenz TJ, Tchong JE. Design and methodology of the ESPRIT trial: evaluating a novel dosing regimen of eptifibatid in percutaneous coronary intervention. *Am Heart J* 2000; **140**: 834–9.
 - 14 O'Shea JC, Madan M, Cantor WJ, Pacchiana CM, Greenberg S, Joseph DM, Kitt MM, Lorenz TJ, Tchong JE. Platelet glycoprotein IIb/IIIa receptor blockade and low-dose heparin during percutaneous coronary revascularization. The EPILOG Investigators. *N Engl J Med* 1997; **336**: 1689–96.
 - 15 O'Shea JC, Madan M, Cantor WJ, Pacchiana CM, Greenberg S, Joseph DM, Kitt MM, Lorenz TJ, Tchong JE. Use of a monoclonal antibody directed against the platelet glycoprotein IIb/IIIa receptor in high-risk coronary angioplasty. The EPIC Investigation. *N Engl J Med* 1994; **330**: 956–61.
 - 16 Cohen M, Demers C, Gurfinkel EP, Turpie AG, Fromell GJ, Goodman S, Langer A, Califf RM, Fox KA, Premeureur J, Bigonzi F. A comparison of low-molecular-weight heparin with unfractionated heparin for unstable coronary artery disease. Efficacy and Safety of Subcutaneous Enoxaparin in Non-Q-Wave Coronary Events Study Group. *N Engl J Med* 1997; **337**: 447–52.
 - 17 Antman EM, McCabe CH, Gurfinkel EP, Turpie AG, Bernink PJ, Salein D, Bayes De Luna A, Fox K, Lablanche JM, Radley D, Premeureur J, Braunwald E. Enoxaparin prevents death and cardiac ischemic events in unstable angina/non-Q-wave myocardial infarction. Results of the thrombolysis in myocardial infarction (TIMI) 11B trial. *Circulation* 1999; **100**: 1593–601.
 - 18 Turpie AG, Bauer KA, Eriksson BI, Lassen MR. Fondaparinux vs enoxaparin for the prevention of venous thromboembolism in major orthopedic surgery: a meta-analysis of 4 randomized double-blind studies. *Arch Intern Med* 2002; **162**: 1833–40.
 - 19 Leon MB, Baim DS, Popma JJ, Gordon PC, Cutlip DE, Ho KK, Giambartolomei A, Diver DJ, Lasorda DM, Williams DO, Pocock SJ, Kuntz RE. A clinical trial comparing three antithrombotic-drug regimens after coronary-artery stenting. Stent Anticoagulation Restenosis Study Investigators. *N Engl J Med* 1998; **339**: 1665–71.
 - 20 Bertrand ME, Legrand V, Boland J, Fleck E, Bonnier J, Emmanuelson H, Vrolix M, Missault L, Chierchia S, Casaccia M, Niccoli L, Oto A, White C, Webb-Peplow M, Van Belle E, McFadden EP. Randomized multicenter comparison of conventional anticoagulation versus antiplatelet therapy in unplanned and elective coronary stenting. The full anticoagulation versus aspirin and ticlopidine (fantastic) study. *Circulation* 1998; **98**: 1597–603.
 - 21 Bhatt DL, Topol EJ. Scientific and therapeutic advances in antiplatelet therapy. *Nat Rev Drug Discov* 2003; **2**: 15–28.
 - 22 Moshfegh K, Redondo M, Julmy F, Wuillemin WA, Gebauer MU, Haeberli A, Meyer BJ. Antiplatelet effects of clopidogrel compared with aspirin after myocardial infarction: enhanced inhibitory effects of combination therapy. *J Am Coll Cardiol* 2000; **36**: 699–705.
 - 23 Andre P, LaRocca T, Delaney SM, Lin PH, Vincent D, Sinha U, Conley PB, Phillips DR. Anticoagulants (thrombin inhibitors) and aspirin synergize with P2Y₁₂ receptor antagonism in thrombosis. *Circulation* 2003; **108**: 2697–703.
 - 24 Lorrain J, Lechaire I, Gauffeny C, Masson R, Roome N, Haurault JP, O'Connor SE, Schaeffer P, Herbert JM. Effects of SanOrg123781A, a synthetic hexadecasaccharide, in a mouse model of electrically induced carotid artery injury: synergism with the antiplatelet agent clopidogrel. *J Pharmacol Exp Ther* 2004; **309**: 235–40.
 - 25 Ni H, Papalia JM, Degen JL, Wagner DD. Control of thrombus embolization and fibronectin internalization by integrin alpha IIb beta 3 engagement of the fibrinogen gamma chain. *Blood* 2003; **102**: 3609–14.
 - 26 Jin J, Kunapuli SP. Coactivation of two different G protein-coupled receptors is essential for ADP-induced platelet aggregation. *Proc Natl Acad Sci USA* 1998; **95**: 8070–4.
 - 27 Moers A, Nieswandt B, Massberg S, Wettschreck N, Gruner S, Konrad I, Schulte V, Aktas B, Gratacap MP, Simon MI, Gawaz M, Offermanns S. G13 is an essential mediator of platelet activation in hemostasis and thrombosis. *Nat Med* 2003; **9**: 1418–22.
 - 28 Turner NA, Moake JL, McIntire LV. Blockade of adenosine diphosphate receptors P2Y₁₂ and P2Y₁ is required to inhibit platelet aggregation in whole blood under flow. *Blood* 2001; **98**: 3340–5.
 - 29 Ferguson JJ, Califf RM, Antman EM, Cohen M, Grines CL, Goodman S, Kereiakes DJ, Langer A, Mahaffey KW, Nessel CC, Armstrong PW, Avezum A, Aylward P, Becker RC, Biasucci L, Borzak S, Col J, Frey MJ, Fry E, Gulba DC, Guneri S, Gurfinkel E, Harrington R, Hochman JS, Kleiman NS, Leon MB, Lopez-Sendon JL, Pepine CJ, Ruzyllo W, Steinhubl SR, Teirstein PS, Toro-Figueroa L, White H. Enoxaparin vs unfractionated heparin in high-risk patients with non-ST-segment elevation acute coronary syndromes managed with an intended early invasive strategy: primary results of the SYNERGY randomized trial. *J Am Med Assoc* 2004; **292**: 45–54.
 - 30 Gum PA, Kottke-Marchant K, Welsh PA, White J, Topol EJ. A prospective, blinded determination of the natural history of aspirin resistance among stable patients with cardiovascular disease. *J Am Coll Cardiol* 2003; **41**: 961–5.
 - 31 Gum PA, Kottke-Marchant K, Poggio ED, Gurm H, Welsh PA, Brooks L, Sapp SK, Topol EJ. Profile and prevalence of aspirin resistance in patients with cardiovascular disease. *Am J Cardiol* 2001; **88**: 230–5.
 - 32 Eikelboom JW, Hirsh J, Weitz JI, Johnston M, Yi Q, Yusuf S. Aspirin-resistant thromboxane synthesis and the risk of myocardial infarction, stroke, or cardiovascular death in patients at high risk for cardiovascular events. *Circulation* 2002; **105**: 1650–5.
 - 33 Zimmermann N, Wenk A, Kim U, Kienzle P, Weber AA, Gams E, Schror K, Hohlfeld T. Functional and biochemical evaluation of platelet aspirin resistance after coronary artery bypass surgery. *Circulation* 2003; **108**: 542–7.
 - 34 Catella-Lawson F, Reilly MP, Kapoor SC, Cucchiara AJ, DeMarco S, Tournier B, Vyas SN, FitzGerald GA. Cyclooxygenase inhibitors and the antiplatelet effects of aspirin. *N Engl J Med* 2001; **345**: 1809–17.

- 35 Topol E, Quinn MJ. Common variations in platelet glycoproteins: pharmacogenomic implications. *Pharmacogenomics* 2001; **2**: 341–52.
- 36 Serebruany VL, Steinhubl SR, Berger PB, Malinin AI, Bhatt DL, Topol EJ. Variability in platelet responsiveness to clopidogrel among 544 individuals. *J Am Coll Cardiol* 2005; **45**: 246–51.
- 37 Taubert D, Kastrati A, Harlfinger S, Gorchakova O, Lazar A, von Beckerath N, Schomig A, Schomig E. Pharmacokinetics of clopidogrel after administration of a high loading dose. *Thromb Haemost* 2004; **92**: 311–6.
- 38 Muller I, Besta F, Schulz C, Massberg S, Schonig A, Gawaz M. Prevalence of clopidogrel non-responders among patients with stable angina pectoris scheduled for elective coronary stent placement. *Thromb Haemost* 2003; **89**: 783–7.
- 39 Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Ramirez C, Sabate M, Banuelos C, Hernandez-Antolin R, Escaned J, Moreno R, Alfonso F, Macaya C. High clopidogrel loading dose during coronary stenting: effects on drug response and interindividual variability. *Eur Heart J* 2004; **25**: 1903–10.
- 40 Gurbel PA, Bliden KP, Hiatt BL, O'Connor CM. Clopidogrel for coronary stenting: response variability, drug resistance, and the effect of pretreatment platelet reactivity. *Circulation* 2003; **107**: 2908–13.
- 41 Matetzky S, Shenkman B, Guetta V, Shechter M, Bienart R, Goldenberg I, Novikov I, Pres H, Savion N, Varon D, Hod H. Clopidogrel resistance is associated with increased risk of recurrent atherothrombotic events in patients with acute myocardial infarction. *Circulation* 2004; **109**: 3171–5.
- 42 Lau WC, Waskell LA, Watkins PB, Neer CJ, Horowitz K, Hopp AS, Tait AR, Carville DG, Guyer KE, Bates ER. Atorvastatin reduces the ability of clopidogrel to inhibit platelet aggregation: a new drug-drug interaction. *Circulation* 2003; **107**: 32–7.
- 43 Blazing MA, de Lemos JA, White HD, Fox KA, Verheugt FW, Ardissino D, DiBattiste PM, Palmisano J, Bilheimer DW, Snapinn SM, Ramsey KE, Gardner LH, Hasselblad V, Pfeffer MA, Lewis EF, Braunwald E, Califf RM. Safety and efficacy of enoxaparin vs unfractionated heparin in patients with non-ST-segment elevation acute coronary syndromes who receive tirofiban and aspirin: a randomized controlled trial. *J Am Med Assoc* 2004; **292**: 55–64.
- 44 Blazing MA, de Lemos JA, White HD, Fox KA, Verheugt FW, Ardissino D, DiBattiste PM, Palmisano J, Bilheimer DW, Snapinn SM, Ramsey KE, Gardner LH, Hasselblad V, Pfeffer MA, Lewis EF, Braunwald E, Califf RM. The Thrombolysis in Myocardial Infarction (TIMI) 11A Trial Investigators. *J Am Coll Cardiol* 1997; **29**: 1474–82.
- 45 Lincoff AM, Bittl JA, Harrington RA, Feit F, Kleiman NS, Jackman JD, Sarembock IJ, Cohen DJ, Spriggs D, Ebrahimi R, Keren G, Carr J, Cohen EA, Betriu A, Desmet W, Kereiakes DJ, Rutsch W, Wilcox RG, De Feyter PJ, Vahanian A, Topol EJ. Bivalirudin and provisional glycoprotein IIb/IIIa blockade compared with heparin and planned glycoprotein IIb/IIIa blockade during percutaneous coronary intervention: REPLACE-2 randomized trial. *J Am Med Assoc* 2003; **289**: 853–63.
- 46 Francis CW, Berkowitz SD, Comp PC, Lieberman JR, Ginsberg JS, Paiement G, Peters GR, Roth AW, McElhattan J, Colwell CW Jr. Comparison of ximelagatran with warfarin for the prevention of venous thromboembolism after total knee replacement. *N Engl J Med* 2003; **349**: 1703–12.
- 47 Olsson SB. Stroke prevention with the oral direct thrombin inhibitor ximelagatran compared with warfarin in patients with non-valvular atrial fibrillation (SPORTIF III): randomised controlled trial. *Lancet* 2003; **362**: 1691–8.
- 48 Damiano BP, Derian CK, Maryanoff BE, Zhang HC, Gordon PA. RWJ-58259: a selective antagonist of protease activated receptor-1. *Cardiovasc Drug Rev* 2003; **21**: 313–26.
- 49 Kawahara TSS, Matsuura F, Clark RSJ et al. Discovery and optimization of potent orally active small molecular thrombin receptor (PAR-1) antagonists. In: *Medi 85. 227 ACS National Meeting*, Anaheim, 2004.
- 50 Storey RF, Oldroyd KG, Wilcox RG. Open multicentre study of the P2_T receptor antagonist AR-C69931MX assessing safety, tolerability and activity in patients with acute coronary syndromes. *Thromb Haemost* 2001; **85**: 401–7.
- 51 Springthorpe B. *From ATP to AZD6140: Design of an Orally Active P2Y₁₂ Receptor Antagonist for the Treatment of Thrombosis*. New Orleans: American Chemical Society Meeting, 2003.
- 52 Springthorpe B. Best of the ESC 2004. *Rev Cardiovasc Med* 2004; **5**: 223–5.
- 53 Hoppelner G, Jantzen HM, Vincent D, Li G, England L, Ramakrishnan V, Yang RB, Nurden P, Nurden A, Julius D, Conley PB. Identification of the platelet ADP receptor targeted by antithrombotic drugs. *Nature* 2001; **409**: 202–7.
- 54 Foster CJ, Prosser DM, Agans JM, Zhai Y, Smith MD, Lachowicz JE, Zhang FL, Gustafson E, Monsma FJ Jr, Wiekowski MT, Abbondanzo SJ, Cook DN, Bayne ML, Lira SA, Chintala MS. Molecular identification and characterization of the platelet ADP receptor targeted by thienopyridine antithrombotic drugs. *J Clin Invest* 2001; **107**: 1591–8.
- 55 Leon C, Hechler B, Freund M, Eckly A, Vial C, Ohlmann P, Dierich A, LeMeur M, Cazenave JP, Gachet C. Defective platelet aggregation and increased resistance to thrombosis in purinergic P2Y₁(1) receptor-null mice. *J Clin Invest* 1999; **104**: 1731–7.
- 56 Fabre JE, Nguyen M, Latour A, Keifer JA, Audoly LP, Coffman TM, Koller BH. Decreased platelet aggregation, increased bleeding time and resistance to thromboembolism in P2Y₁-deficient mice. *Nat Med* 1999; **5**: 1199–202.
- 57 Andre P, Delaney SM, LaRocca T, Vincent D, DeGuzman F, Jurek M, Koller B, Phillips DR, Conley PB. P2Y₁₂ regulates platelet adhesion/activation, thrombus growth, and thrombus stability in injured arteries. *J Clin Invest* 2003; **112**: 398–406.
- 58 Kim HS, Ohno M, Xu B, Kim HO, Choi Y, Ji XD, Maddileti S, Marquez VE, Harden TK, Jacobson KA. 2-Substitution of adenine nucleotide analogues containing a bicyclo[3.1.0]hexane ring system locked in a northern conformation: enhanced potency as P2Y₁ receptor antagonists. *J Med Chem* 2003; **46**: 4974–87.
- 59 Mathieu R, Baurand A, Schmitt M, Gachet C, Bourguignon JJ. Synthesis and biological activity of 2-alkylated deoxyadenosine bisphosphate derivatives as P2Y₁ receptor antagonists. *Bioorg Med Chem* 2004; **12**: 1769–79.
- 60 Law DA, DeGuzman FR, Heiser P, Ministri-Madrid K, Killeen N, Phillips DR. Integrin cytoplasmic tyrosine motif is required for outside-in alphaIIb beta3 signalling and platelet function. *Nature* 1999; **401**: 808–11.
- 61 Andre P, Nannizzi-Alaimo L, Prasad SK, Phillips DR. Platelet-derived CD40L: the switch-hitting player of cardiovascular disease. *Circulation* 2002; **106**: 896–9.
- 62 Andre P, Prasad KS, Denis CV, He M, Papalia JM, Hynes RO, Phillips DR, Wagner DD. CD40L stabilizes arterial thrombi by a beta3 integrin-dependent mechanism. *Nat Med* 2002; **8**: 247–52.
- 63 Prasad KS, Andre P, Yan Y, Phillips DR. The platelet CD40L/GP IIb-IIIa axis in atherothrombotic disease. *Curr Opin Hematol* 2003; **10**: 356–61.
- 64 Angelillo-Scherrer A, de Frutos P, Aparicio C, Melis E, Savi P, Lupu F, Arnout J, Dewerchin M, Hoylaerts M, Herbert J, Collen D, Dahlback B, Carmeliet P. Deficiency or inhibition of Gas6 causes platelet dysfunction and protects mice against thrombosis. *Nat Med* 2001; **7**: 215–21.
- 65 Angelillo-Scherrer A, Burnier L, Flores N, Savi P, Demol M, Schaeffer P, Herbert JM, Lemke G, Goff SP, Matsushima GK, Earp HS, Vesin C, Hoylaerts MF, Plaisance S, Collen D, Conway EM, Wehrle-Haller B, Carmeliet P. Role of Gas6 receptors in platelet signaling during thrombus stabilization and implications for antithrombotic therapy. *J Clin Invest* 2005; **115**: 237–46.
- 66 Prevost N, Woulfe D, Tanaka T, Brass LF. Interactions between Eph kinases and ephrins provide a mechanism to support platelet aggregation.

- gation once cell-to-cell contact has occurred. *Proc Natl Acad Sci USA* 2002; **99**: 9219–24.
- 67 Siljander PR, Munnix IC, Smethurst PA, Deckmyn H, Lindhout T, Ouwehand WH, Farndale RW, Heemskerk JW. Platelet receptor interplay regulates collagen-induced thrombus formation in flowing human blood. *Blood* 2004; **103**: 1333–41.
- 68 Moroi M, Jung SM, Shinmyozu K, Tomiyama Y, Ordinas A, Diaz-Ricart M. Analysis of platelet adhesion to a collagen-coated surface under flow conditions: the involvement of glycoprotein VI in the platelet adhesion. *Blood* 1996; **88**: 2081–92.
- 69 Bellucci S, Huisse MG, Boval B, Hainaud P, Robert A, Fauvel-Lafeve F, Jandrot-Perrus M. Defective collagen-induced platelet activation in two patients with malignant haemopathies is related to a defect in the GPVI-coupled signalling pathway. *Thromb Haemost* 2005; **93**: 130–8.
- 70 Monnet E, Fauvel-Lafeve F. A new platelet receptor specific to type III collagen. Type III collagen-binding protein. *J Biol Chem* 2000; **275**: 10912–7.
- 71 Moroi M, Jung SM, Okuma M, Shinmyozu K. A patient with platelets deficient in glycoprotein VI that lack both collagen-induced aggregation and adhesion. *J Clin Invest* 1989; **84**: 1440–5.
- 72 Arai M, Yamamoto N, Moroi M, Akamatsu N, Fukutake K, Tanoue K. Platelets with 10% of the normal amount of glycoprotein VI have an impaired response to collagen that results in a mild bleeding tendency. *Br J Haematol* 1995; **89**: 124–30.
- 73 Jackson SP, Yap CL, Anderson KE. Phosphoinositide 3-kinases and the regulation of platelet function. *Biochem Soc Trans* 2004; **32**: 387–92.
- 74 Offermanns S, Toombs CF, Hu YH, Simon MI. Defective platelet activation in G alpha(q)-deficient mice. *Nature* 1997; **389**: 183–6.
- 75 Jantzen HM, Milstone DS, Gousset L, Conley PB, Mortensen RM. Impaired activation of murine platelets lacking G α_{i2} . *J Clin Invest* 2001; **108**: 477–83.
- 76 Yang J, Wu J, Jiang H, Mortensen R, Austin S, Manning DR, Woulfe D, Brass LF. Signaling through Gi family members in platelets. Redundancy and specificity in the regulation of adenylyl cyclase and other effectors. *J Biol Chem* 2002; **277**: 46035–42.
- 77 Obergfell A, Eto K, Mocsai A, Buensuceso C, Moores SL, Brugge JS, Lowell CA, Shattil SJ. Coordinate interactions of Csk, Src, and Syk kinases with [alpha]IIb[beta]3 initiate integrin signaling to the cytoskeleton. *J Cell Biol* 2002; **157**: 265–75.
- 78 Law DA, Nannizzi-Alaimo L, Ministri K, Hughes PE, Forsyth J, Turner M, Shattil SJ, Ginsberg MH, Tybulewicz VL, Phillips DR. Genetic and pharmacological analyses of Syk function in alphaIIb-beta3 signaling in platelets. *Blood* 1999; **93**: 2645–52.
- 79 Judd BA, Myung PS, Leng L, Obergfell A, Pear WS, Shattil SJ, Koretzky GA. Hematopoietic reconstitution of SLP-76 corrects hemostasis and platelet signaling through alpha IIb beta 3 and collagen receptors. *Proc Natl Acad Sci USA* 2000; **97**: 12056–61.
- 80 Schober A, Manka D, von Hundelshausen P, Huo Y, Hanrath P, Sarembock IJ, Ley K, Weber C. Deposition of platelet RANTES triggering monocyte recruitment requires P-selectin and is involved in neointima formation after arterial injury. *Circulation* 2002; **106**: 1523–9.
- 81 Huo Y, Schober A, Forlow SB, Smith DF, Hyman MC, Jung S, Littman DR, Weber C, Ley K. Circulating activated platelets exacerbate atherosclerosis in mice deficient in apolipoprotein E. *Nat Med* 2003; **9**: 61–7.
- 82 Burger PC, Wagner DD. Platelet P-selectin facilitates atherosclerotic lesion development. *Blood* 2003; **101**: 2661–6.
- 83 Ault KA, Cannon CP, Mitchell J, McCahan J, Tracy RP, Novotny WF, Reimann JD, Braunwald E. Platelet activation in patients after an acute coronary syndrome: results from the TIMI-12 trial. Thrombolysis in myocardial infarction. *J Am Coll Cardiol* 1999; **33**: 634–9.
- 84 May AE, Neumann FJ, Gawaz M, Ott I, Walter H, Schomig A. Reduction of monocyte-platelet interaction and monocyte activation in patients receiving antiplatelet therapy after coronary stent implantation. *Eur Heart J* 1997; **18**: 1913–20.
- 85 Mach F, Schonbeck U, Sukhova GK, Atkinson E, Libby P. Reduction of atherosclerosis in mice by inhibition of CD40 signalling. *Nature* 1998; **394**: 200–3.
- 86 Lutgens E, Gorelik L, Daemen MJ, de Muinck ED, Grewal IS, Kotliansky VE, Flavell RA. Requirement for CD154 in the progression of atherosclerosis. *Nat Med* 1999; **5**: 1313–6.
- 87 Heeschen C, Dimmeler S, Hamm CW, van den Brand MJ, Boersma E, Zeiher AM, Simoons ML. Soluble CD40 ligand in acute coronary syndromes. *N Engl J Med* 2003; **348**: 1104–11.
- 88 Quinn MJ, Bhatt DL, Zidar F, Vivekananthan D, Chew DP, Ellis SG, Plow E, Topol EJ. Effect of clopidogrel pretreatment on inflammatory marker expression in patients undergoing percutaneous coronary intervention. *Am J Cardiol* 2004; **93**: 679–84.
- 89 Massberg S, Brand K, Gruner S, Page S, Muller E, Muller I, Bergmeier W, Richter T, Lorenz M, Konrad I, Nieswandt B, Gawaz M. A critical role of platelet adhesion in the initiation of atherosclerotic lesion formation. *J Exp Med* 2002; **196**: 887–96.
- 90 Lynch TJ, Bell DW, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, Harris PL, Haserlat SM, Supko JG, Haluska FG, Louis DN, Christiani DC, Settleman J, Haber DA. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004; **350**: 2129–39.
- 91 Phillips DR, Teng W, Arfsten A, Nannizzi-Alaimo L, White MM, Longhurst C, Shattil SJ, Randolph A, Jakubowski JA, Jennings LK, Scarborough RM. Effect of Ca $^{2+}$ on GP IIb-IIIa interactions with integrilin: enhanced GP IIb-IIIa binding and inhibition of platelet aggregation by reductions in the concentration of ionized calcium in plasma anticoagulated with citrate. *Circulation* 1997; **96**: 1488–94.
- 92 Sakariassen KS, Turitto VT, Baumgartner HR. Recollections of the development of flow devices for studying mechanisms of hemostasis and thrombosis in flowing whole blood. *J Thromb Haemost* 2004; **2**: 1681–90.