



A 34-year-old woman with a 3-year history of systemic lupus erythematosus was admitted to the hospital with sore throat and headache that had lasted for 3 weeks. She had been receiving cyclophosphamide for lupus nephritis for the previous 2 years. On the eighth hospital day, she reported severe headache and bilateral pain and bruising of the eyes. A day later, proptosis developed, with severe chemosis and eyelid edema, and her pupils were bilaterally nonreactive to light on examination.

## IMAGES IN CLINICAL MEDICINE

## Cavernous Sinus Thrombosis



Ayaz Agayev, M.D.  
Sabri Yilmaz, M.D.

Istanbul Faculty of Medicine  
Istanbul, Turkey  
ayazagayev@gmail.com

**A** 34-YEAR-OLD WOMAN WITH A 3-YEAR HISTORY OF SYSTEMIC LUPUS ERYTHEMATOSUS was admitted to the hospital with sore throat and headache that had lasted for 3 weeks. She had been receiving cyclophosphamide for lupus nephritis for the previous 2 years. On the eighth hospital day, she reported severe headache and bilateral pain and bruising of the eyes. A day later, proptosis developed, with severe chemosis and eyelid edema, and her pupils were bilaterally non-reactive to light on examination. A preliminary diagnosis of extensive intracranial venous thrombosis was confirmed on magnetic resonance imaging. Axial T<sub>2</sub>-weighted imaging at the level of the cavernous sinus was performed. As compared with an image from a healthy adult (Fig. 1A in the Supplementary Appendix, available with the full text of this article at [www.nejm.org](http://www.nejm.org)), the patient's image showed enlargement and heterogeneity in bilateral cavernous sinuses and in the right transverse sinus, with loss of flow void (Fig. 1B in the Supplementary Appendix). Bilateral exophthalmos, thickening of extraocular muscles, periorbital edema, and hyperintense signal compatible with inflammation of the bilateral mastoid cells (Fig. 1B in the Supplementary Appendix) were also observed. Despite treatment with heparin for 5 days, the patient died from cardiac arrest following brain edema.

Copyright © 2008 Massachusetts Medical Society.

## WEB-ONLY IMAGES IN CLINICAL MEDICINE

Click on "Featured Images" at [www.nejm.org](http://www.nejm.org) to see the *Journal's* Web-only Images in Clinical Medicine. The Images are listed (with e page numbers) in the table of contents of the printed *Journal* the week they are published and are compiled on the *Journal's* Web site.



**Supplemental Figure 1.** Axial T<sub>2</sub>-weighted imaging at the cavernous sinus level was performed in a healthy adult (Panel A) and the patient (Panel B). The patient had enlargement and heterogeneity bilateral cavernous sinuses (arrowheads) and in the right transverse sinus (asterisk), with loss of flow void. Bilateral exophthalmos, thickening of extraocular muscles, periorbital edema, and hyperintense signal compatible with inflammation of the bilateral mastoid cells (arrows) were also observed.